Hemorragia intraventricular traumática

Este tipo de hemorragia intraventricular, se presenta en un 10 % de los TCE graves e indican un mal pronóstico [1)].

La hemorragia intraventricular traumática suele asociarse a otras lesiones como contusión intracerebral,hemorragia cerebral, hemorragia subaracnoidea, y edema cerebral difuso.

Suele tener un mal pronóstico.

La mayoría de los pacientes tienen un Glasgow GCS inferior a 8 debido a las lesiones asociadas y la gravedad de la lesión.

La prevalencia es de 0,4 a 4% en todos los pacientes con TCE craneal a los cuales se les practica un TAC cerebral (Is y col., 2012).

Bibliografía

Is, Merih, Ferruh Gezen, Mehmet Akgul, and Murat Dosoglu. 2011. “Traumatic Intraventricular Hemorrhage with a Good Prognosis.” Turkish Neurosurgery 21 (1) (January): 107–109.

1)
Atzema, C., W. R. Mower, et al. (2006). “Prevalence and prognosis of traumatic intraventricular hemorrhage in patients with blunt head trauma.” J Trauma 60(5): 1010-7; discussion 1017.

From:
http://neurocirugiacontemporanea.com/ - **Neurocirugía Contemporánea ISSN 1988-2661**

Permanent link:
http://neurocirugiacontemporanea.com/doku.php?id=hemorragia_intraventricular_traumatica

Last update: **2019/09/26 22:18**